



Improvement of cytotoxicity of mitoxantrone and daunorubicin by candidone, tephrosin, and bavachinin

Sina Darzi^{1,2} · Seyed Abbas Mirzaei¹ · Fatemeh Elahian¹ · Amir Peymani³ · Babak Rahmani⁴ · Shaghayegh Pishkhan Dibazar⁵ · Sadegh Shirian^{6,7} · Leila Shakeri Chaleshtori⁸ · Ehsan Aali³

Received: 5 November 2020 / Accepted: 6 September 2021
© The Author(s), under exclusive licence to Springer Nature B.V. 2021

Abstract

Background Flavonoids have been demonstrated to have the ability of sensitizing cancer cells to chemotherapy and inverse multidrug resistance via various mechanisms, such as modulating of pumps. The therapeutic effect of candidone, tephrosin, and bavachinin in treatment of cancer, particularly to overcome multidrug resistance (MDR) is largely unknown. The capacity of these agents in sensitization of MDR cells is investigated in the current work.

Methods and results We analyzed the impact of candidone, tephrosin, and bavachinin, as chemosensitizer on cell cytotoxicity, *P-gp* and *ABCG2* mRNA expression level on two multidrug resistant cells, *ABCG2* overexpressing human epithelial breast cancer cell line (MCF7/MX), and *P-gp* overexpressing human gastric adenocarcinoma cell line (EPG85.257RDB). The inhibitory concentration of 50% (IC₅₀) of daunorubicin in EPG85.257RDB cells in combination with IC₁₀ of Bavachinin, Tephrosin, and Candidone were 6159 ± 948 , 4186 ± 665 , 730 ± 258 nM, and this data in MCF7/MX cell were 1773 ± 534 , 7160 ± 405 and 3340 ± 622 nM respectively. These three flavonoids dose-dependently decreased the viability of MCF7/MX and EPG85.257RDB and significantly ($p < 0.05$) decreased IC₅₀ of daunorubicin and mitoxantrone except Tephrosin in MCF7/MX cells. Candidone and Bavachinin were the most potent chemosensitizer in EPG85.257RDB and MCF7/MX cells respectively. Flavonoids did not reduce mRNA expression of *P-gp* and *ABCG2* after 72 h treatment, except Candidone in EPG85.257RDB and Bavachinin in MCF7/MX cells.

Conclusions This effect is not time-dependent, and flavonoids have their own patterns that are cell-dependent. In general, tephrosin, candidone, and bavachinin had the potential of sensitizing MDR cells to mitoxantrone and daunorubicin.

Keywords MDR1 · *ABCG2* · Flavonoid · Multidrug resistant · Real-time pcr

✉ Ehsan Aali
en.aali@gmail.com; e.aali@qums.ac.ir

Sina Darzi
S.darzi@qums.ac.ir

Seyed Abbas Mirzaei
mirzaei.a@skums.ac.ir

Fatemeh Elahian
elahian.f@skums.ac.ir

Amir Peymani
apeymani@qums.ac.ir

Babak Rahmani
brahmani@qums.ac.ir

Sadegh Shirian
shirian85@gmail.com

² Health Product Safety Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

³ Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

⁴ Department of Molecular Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

⁵ Department of Biotechnology, Qazvin University of Medical Sciences, Qazvin, Iran

⁶ Department of Pathology, School of Veterinary Medicine, Shahrekord University, Shahrekord, Iran

⁷ Shiraz Molecular Research Center, Dr. Daneshbod Lab, Shiraz, Iran

⁸ Department of Biology, Faculty of Basic Sciences, Payame Noor University (PNU), Isfahan, Iran

¹ Department of Molecular Medicine, Faculty of Advanced Technologies, Shahrekord University of Medical Sciences, Shahrekord, Iran